

## A preliminary survey of amphibians from the Idanre Forest Ecosystem, southwestern Nigeria

Angela Ijeoma Ugo<sup>1</sup>, Lotanna Micah Nneji<sup>2\*</sup>, Babatunde Emmanuel Adedeji<sup>1</sup>, Akindele Oluwatosin Adeyi<sup>3\*</sup>

<sup>1</sup>Ecology and Environmental Biology Unit, Department of Zoology, University of Ibadan, Ibadan, Oyo State, Nigeria

<sup>2</sup>Herpetological Diversity and Evolution Unit, State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, PR China

<sup>3</sup>Animal Physiology Unit, Department of Zoology, University of Ibadan, Ibadan, Oyo State, Nigeria \*Email: lotannanncji@gmail.com; delegenius@yahoo.com

Received: 13 November 2020 / Revised: 05 April 2021 / Accepted: 05 April 2021 / Published online: 05 April 2021. Ministry of Sciences, Research, and Technology, Arak University, Iran.

**How to cite:** Ugo, A.I., Nneji, L.M., Adedeji, B.E., Adeyi, A.O. (2021). A preliminary survey of amphibians from the Idanre Forest Ecosystem, southwestern Nigeria. *Journal of Wildlife and Biodiversity*, 5 (4), 1-9. [DOI:10.22120/jwb.2021.139825.1201](https://doi.org/10.22120/jwb.2021.139825.1201)


### Abstract

This study was conducted to provide a checklist of amphibian species in the Idanre forest ecosystem (lowland - Idanre Forest Reserve, and highland - Idanre Hills) based on field surveys carried out during June - July 2018. A total of 218 individuals, comprising 11 species from six genera belonging to five families, were observed. A high number of individuals and species were observed in the lowland (Idanre Forest Reserve) compared to the highland (Idanre Hills) region. All species recorded from the study sites are categorized as Least Concern in the IUCN Red List, except for the critically endangered and endemic *Sclerophrys perreti*. We acknowledge that this study is preliminary; thus, we offer recommendations for a comprehensive survey to unveil the amphibian species richness and abundance as well as patterns of species' adaptation.

**Keywords:** Diversity, species richness; conservation; amphibian; Nigeria

### Introduction

Amphibians include the highest number of threatened species than other vertebrate groups in the world (Howard & Bickford, 2014). Currently, the International Union for Conservation of Nature (IUCN) reported that approximately one-third of the world's amphibian species are threatened. Among these threats, habitat alteration has been opined as a leading cause of amphibian decline globally (Stuart et al., 2004; Wake & Vredenburg, 2008; Hof et al., 2011). Other threats include invasive species, disease, and climate change (Cheng et al., 2011; Hof et al., 2011)



**Figure 1.** Map of Nigeria, showing the Idanre forest ecosystem, Ondo State, southwestern Nigeria. Owing to the global decline of amphibians, there is a need for improved conservation of amphibians and their associated habitats (Nneji *et al.*, 2019a).

Studies on amphibians of Nigeria dates to the 1950s (Romer, 1953; Schiøtz, 1963, 1966; Walker, 1966, 1967, 1968, 1969). In recent times, there has been an increase in amphibian surveys across the country (Akani & Luiselli, 2009; Onadeko & Rödel 2009; Akani *et al.*, 2014; Nneji *et al.*, 2019a, b, c & d, and others). Such studies have continued to give deep insights into the distribution, diversity, and conservation status of amphibians in Nigeria. Extensive herpetological surveys will help understand the amphibian diversity and enable the design of sustainable conservation plans for amphibians and their associated habitats in Nigeria.

Idanre forest ecosystem, consisting of Idanre Forest Reserve and Idanre Hills in its lowland and highland regions respectively, contains a sizeable amount of plant and animal species, some of which are endemic. For instance, previous studies of Onadeko and Rodel (2014) and Nneji *et al.* (2019d) reported an endemic and critically endangered toad – *Sclerophrys perreti* (Schiøtz, 1963) from the Idanre Hills. To date, there is still no comprehensive information on the amphibian species richness in the lowland Idanre forest reserve and its adjacent Idanre Hills. The recent study of Awoku and Ogunjemite (2019) reported a reduction of Nigerian forest reserves' landmass due to the demand for forest land for agriculture, fuelwood, timbers, etc. The apparent human-induced activities in the forest ecosystem in Nigeria, such as hunting, slash-and-burn agriculture, urbanization, etc., pose severe threats to wildlife and its habitats (Ikemeh, 2013; Nneji *et al.*, 2019a). These conservation challenges heighten the need for the documentation of amphibians associated with the Idanre forest ecosystem. Herein, we present the results of our preliminary surveys of amphibians in the Idanre forest ecosystem. The objectives of our study are to (1) create a checklist of amphibians, and (2) compare the amphibian species richness in the lowland and highland areas of the Idanre forest ecosystem.

## Materials and methods

### Study site

Our study area (Fig.1) – the Idanre forest ecosystem – encompasses Idanre Forest Reserve and Idanre Hills. It is located in Ondo State, Southwestern Nigeria. The Idanre Forest reserve is a lowland forest with an altitude of 10 - 400 meters above sea level. It is located between longitude 04°59' E to 5°12' E and latitude 6°45' N to 6°58' N. On the other hand, the Idanre Hills is located between longitude 5°00' E to 5°15' E and latitude 7°00' N to 7°15' N. It resides 3000 ft above sea level and it is characterized by valleys

interspersed by the inselberg and hilly forest across the area. The mean annual rainfall of the Idanre forest ecosystem is about 1,60-2000 mm (Ige *et al.*, 2011; Orimaye *et al.*, 2012) with July and September considered being the period of highest rainfall. The dry season usually lasts for about 3-4 months spanning November through February. The average temperature of the area is between 24 – 34°C (Ige *et al.*, 2011).

### Specimen collection, identification, and diversity analyses

A preliminary survey was carried out during the rainy season for 28 days from 2nd July 2018 to 30th July 2018. Diurnal (05:00 - 08:00) and nocturnal (18:00 - 22:00) searches were conducted using the visual encounter and opportunistic survey method (Rödel and Ernst, 2004). Different microhabitats, including streams, under rocks, within crevices, forest, agricultural farmland, and under wet logs (Fig. 2), were searched randomly for amphibians. Voucher specimens were euthanized with hydrous chlorobutanol, and tissue (muscles) samples were collected and subsequently stored in 95% ethanol for DNA-based identification.



**Figure 2.** Picture showing some of the sampled localities within Idanre Hills, Southwestern Nigeria.

The voucher specimens were fixed with 10% formalin and preserved in 75% ethanol. Specimens were deposited in the museum of the Department of Zoology, University of Ibadan, Nigeria. Preliminary identification of the specimen was conducted using primary literature (Rödel, 2000). For the DNA-based identification, the phenol-chloroform method by Sambrook *et al.* (1989) was followed. A partial mitochondrial 16S ribosomal RNA was amplified and used for genetic identification. The primer pairs designed by Bossuyt and Milinkovitch (2000) were used for the amplification. The molecular-based identification followed the methods of Nneji *et al.* (2019c). Further, we referred to the IUCN Red List to evaluate the threat status of the species (IUCN, 2020). We estimated the relative abundance of each of the species encountered.

### Results and discussion

A total of 218 individuals comprising 11 species distributed in five families of the order Anura were observed (Table 1). No record of caecilian (order: Gymnophiona) was observed from the area. Suffice to say that there is a high likelihood of encountering more amphibian species from the Idanre forest

ecosystem. However, factors such as area restriction due to traditional belief at Idanre Hills, inaccessibility of some areas due to rainfall, sampling techniques, and short duration of field sampling precluded an extensive survey of the area. Thus, we recommend a continuous field survey to document the amphibian compositions of the Idanre forest ecosystem.

Family Bufonidae and Ptychadenidae had the highest number of species representatives ( $n = 3$  species). The most abundant species were *Phrynobatrachus plicatus* (Günther, 1858) ( $n = 57$ ;  $r = 26.20\%$ ) and *Ptychadena longirostris* (Peters, 1870) ( $n = 45$ ;  $r = 20.60\%$ ) (Table 1). Species such as *Leptopelis aubryi* (Duméril, 1856) and *Ptychadena bibroni* (Hallowell, 1845) were encountered less frequently ( $n = 3$ ;  $r = 1.40\%$ ) (Table 1). The IUCN Red List showed ten species from the Idanre forest ecosystem as Least Concern and one (*S. perreti*) as Critically Endangered (Table 1). Previously, *S. perreti*, endemic to the Idanre Hills, was declared lost. However, studies by Onadeko et al. (2014) and Nneji et al. (2019d) reconfirmed its presence and described the species as micro endemic. During our field survey, we observed *S. perreti* from batholithic granite rocks and also within low herbal vegetation that grows on the soil between the rocks. The varied skin coloration of *S. perreti* provides a perfect camouflage to this microhabitat and may have accounted for its persistence in the ecosystem. Comparing both study sites, result showed that the species richness is high ( $n = 9$  species,  $n = 184$  individuals;  $r = 84.40\%$ ) in the lowland region compared to the highland region ( $n = 5$  species,  $n = 34$  individuals;  $r = 15.60\%$ ). Although a bias-related sampling procedure (382 person-hours in the lowland compared to 153 person-hour at the highland) may contribute to this finding, we most likely attribute the abundance of amphibians in the lowland region to the presence of suitable environmental conditions such as humid forest, swamps, leaf-litters, etc. The Idanre Hills comprise rock-outcrop habitats with direct sun exposure and this might have accounted for the fewer number of amphibians observed from the area.

Our survey revealed three species (*Arthroleptis poecilinotus* (Peters, 1863) *Hoplobatrachus occipitalis* (Günther, 1858), and *Sclerophrys regularis* (Reuss, 1833)) common to the highland and the lowland forest regions. Previous studies have reported these species from a wide range of habitats. For instance, *A. poecilinotus* has been reported in secondary forests, herbaceous vegetation, and agricultural land. *Hoplobatrachus occipitalis* is also found in dry savannah to highly degraded forests, including highly polluted streams (Nneji et al., 2019a). On the other hand, *S. regularis* has also been reported from the dry savannah region, montane region, agricultural land, and degraded habitats (Nneji et al., 2019a). The ability to adapt to a wide variety of habitats can explain the presence in both study sites.

Potential threats to the amphibians in the Idanre forest ecosystem have been identified anecdotally during the field survey (Fig. 3). These threats include forest fires, a continuous harvest of trees for firewood, extensive use of chemicals in agriculture, slash-and-burn agricultural practices, and forest clearing. Human-induced forest fires and logging are widespread during the farming season, which destroys the habitats of amphibians. Indeed, the direct and indirect effects of these activities could significantly impact the amphibians and their associated habitats in the Idanre forest ecosystem. Our preliminary study demonstrates that the Idanre forest ecosystem has the potential to be an important natural area in terms of amphibian diversity and raises the conservation needs of amphibians and their associated habitats. We recommend a conservation action plan that would involve the government and local community geared towards improving the long-term persistence of the flora and fauna of the Idanre forest ecosystem.



**Figure 3.** Picture showing human activities within Idanre Hills, Southwestern Nigeria

### Acknowledgment

This study was financed in part by the Mohammed bin Zayed Species Conservation Grant (172517120; L.M.N). We thank the Ministry of Environment, Ondo state, Nigeria for providing ethical approval and permit for conducting the field survey in the Idanre forest ecosystem. We also thank Godwin Chinemerem for assisting us during the field survey.

### References

- Akani, G.C. & Luiselli, L. (2009). Aspects of community ecology of amphibians and reptiles at Bonny Island (Nigeria), an area of priority relevance for the petrochemical industry. *African Journal of Ecology*, 48(4), 939 – 948.
- Akani, G.C., Aifesehi, P.E., Petrozzi, F., Amadi, N. & Luiselli, L. (2014). Preliminary surveys of the terrestrial vertebrate fauna (mammals, reptiles, and amphibians) of the Edumanon Forest Reserve, Nigeria. *Tropical Zoology*, 27, 63 – 72.
- Awoku, G.S. & Ogunjemite, B.G. (2019). Preliminary assessment of the distribution of large mammals within Idanre forest reserve, Ondo State, Nigeria. *Journal of Research in Forestry, Wildlife & Environment*, 11(4), 1 – 5.
- Bossuyt, F. & Milinkovitch, M.C. (2000). Convergent adaptive radiations in Madagascan and Asian ranid frogs reveal covariation between larval and adult traits. *Proceedings of the National Academy of Sciences of the United States of America*, 97, 6585 – 6590.
- Cheng, T.L., Rovito, S.M., Wake, D.B. & Vredenburg, V.T. (2011). Coincident mass extirpation of neotropical amphibians with the emergence of the infection fungal pathogen *Batrachochytrium dendrobatidis*. *Proceedings of the National Academy of Sciences of the United States of America*, 108, 9502 – 9507.
- Hof, C., Araujo, M.B., Jetz, W. & Rahbek, C. (2011) Additive threats from pathogens, climate and land-use change for global amphibian diversity. *Nature*, 480, 516 – 519.
- Howard, S.D. & Bickford, D. P. (2014). Amphibians over the edge: silent extinction risk of Data Deficient species. *Diversity and Distributions*, 20, 837 – 846.
- Ige, O., Adeyemi, C., Ogunfolakan, A., Ayansola, A., Olayemi, A., Taiwo, Y., Olayiwola M. & Oyelade, J. (2011). An Inventory of the Geological, Biological and Cultural Resources on Ufe-Oke Hill, Idanre, Southwestern Nigeria. *Natural Resources*, 2, 180 – 190.
- Ikemeh, R.A. (2013). Sustainable forest management in a human-dominated landscape and its implications for biodiversity conservation: a Nigerian lowland forest perspective. Dove press, 9 – 23.
- IUCN - International Union for Conservation of Nature (2020). The IUCN Red List of Threatened Species. Version (2020)-1. <https://www.iucnredlist.org>. Accessed 1 Nov 2020.
- Nneji, L.M., Adeola, A., Okeyoyin, A., Onadeko, A., Eniang, E., Adeyi, O., Oladipo, S., Adedeji, B., Usongo, J., Andem, A., Ugwumba, A., Ugwumba, A., & Olory, C. (2019b). First record of Foulassi Screeching Frog, *Arthroleptis adelphus* (Perret, 1966) (Anura, Arthroleptidae, Arthroleptinae), from Nigeria, with notes on its phylogenetic position. *Check List*, 15 (2), 253 – 258.

- Nneji, L.M., Adeola, A.C., Okeyoyin, A.O., Oladipo, O.C., Saidu, Y., Samuel D, Usongo J.Y., Babatunde A., Olatunde O., Adeyi, A. O., Ugwumba, O.A., Ugwumba, A.A.A (2019a). Diversity and distribution of amphibians and reptiles in Gashaka Gumti National Park, Nigeria. *Herpetology Notes*, 12, 543 – 599.
- Nneji, L.M., Adeola, A.C., Okeyoyin, A.O., Oladipo, O.C., Saidu, Y., Usongo, J.Y., & Ugwumba A.A.A. (2019c). Assessing the effectiveness of molecular data for species identification and diversity studies of Nigerian herpetofauna. *Mitochondrial DNA B*, 4, 589 – 593.
- Nneji, L.M., Salako, G., Adeola, A.C., Oladipo, S.O, Adedeji, B.E., & Ugwumba, A. A. A. (2019d). Species Distribution Modelling predicts habitat suitability and reduction of suitable habitat under future climatic scenario for *Sclerophrys perreti* - a critically endangered Nigerian endemic toad. *African Journal of Ecology*, 00, 1 – 11.
- Onadoko, A., Rödel, M., Liedtke, H., & Barej, M. (2014). The rediscovery of Perret's toad, *Amietophrynus perreti* (Schjøtz, 1963) after more than 40 years, with comments on the species' phylogenetic placement and conservation status. *Zoosystematics and Evolution*, 90(2), 113 – 19.
- Onadoko, A.O. & Rödel, M.O. (2009). Anuran surveys in south–western Nigeria. *Salamandra*, 45 (1), 1 – 14.
- Orimaye, J.O., Ogunjemite, B.G. & Agbelusi, A.E. (2012). Density and Abundance of the Red-Capped Mangabey (*Cercocebus Torquatus*) In Omo Biosphere Reserve and Idanre Forest Reserve, South-Western Nigeria. *MOJ Proteomics & Bioinformatics*, 5 (2), 63 – 67.
- Rödel, M.O. (2000). Herpetofauna of West Africa, Volume I: Amphibians of the West African savannah. Frankfurt am Main, Edition Chimaira.
- Rödel, M.O. & Ernst, R. (2004). Measuring and monitoring amphibian diversity in tropical forests: An evaluation of methods with recommendations for standardization. *Ecotropica*, 10, 1 – 14.
- Romer, J.D. (1953). Reptiles and amphibians were collected in the Port Harcourt area of Nigeria. *Copeia*, 2, 121 – 123.
- Sambrook, J., Fritschi, E.F. & Maniatis, T. (1989). *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, New York.
- Schjøtz, A. (1963). The amphibians of Nigeria. *meddelelser fra Dansk Naturhistorisk Forening*, 125, 1 – 92.
- Schjøtz, A. (1966). On a collection of amphibians from Nigeria. *Videnskabelig meddelelser fra Dansk Naturhistorisk Forening*, 29, 43 – 48.
- Stuart, S.N., Chanson, J.S., Cox, N.A., Young, B.E., Rodrigues, A.S., Fischman, D.L. & Waller, R.W. (2004). Status and trends of amphibian declines and extinctions worldwide. *Science*, 306, 1783 – 1786.
- Wake, D.B. & Vredenburg, V.T. (2008). Are we in the midst of the sixth mass extinction? A review from the world of amphibians. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 11466 – 11473.
- Walker, R.B. (1966). An uncommon frog, *Pyxicephalus adspersus*. *Nigerian Field*, 31, 177–180.
- Walker, R.B. (1967). *Rana galamensis*, an elusive frog. *Nigerian Field*, 32(1), 22 – 26.
- Walker, R.B. (1968). The amphibians of Zaria, in the northern guinea savannah, Nigeria. *Copeia*, 1, 164–167.
- Walker, R.B. (1969). The burrowing frog: *Hemisis marmoratus* (Peters). *Nigerian Field*, 34,72–76.

**Table 1.** Checklist of Amphibian species in Idanre Forest Ecosystem, Ondo State, Southwestern Nigeria

Family	Genus	Species	IUCN status	The total number of individuals observed	Idanre Forest Ecosystem	Idanre Hills
<b>Arthroleptidae</b>						
	<i>Arthroleptis</i>					
		<i>A. poecilinotus</i> (Peters, 1863)	LC	22 (10.10%)	21 (11.40%)	1 (2.90%)
	<i>Leptopelis</i>					
		<i>L. aubryi</i> (Duméril, 1856)	LC	3 (1.40%)	3 (1.60%)	0 (0.00%)
<b>Bufo</b>						
	Sclerophrys					
		<i>S. regularis</i> (Reuss, 1833)	LC	10 (4.60%)	9 (4.90%)	1 (2.90%)
		<i>S. maculata</i> (Hallowell, 1854)	LC	7 (3.20%)	0 (0.00%)	7 (20.60%)
		<i>S. perretti</i> (Schlötter, 1963)	CE	23 (10.60%)	0 (0.00%)	23 (67.70%)
<b>Dicroglossidae</b>						
	<i>Hoplobatrachus</i>	<i>H. occipitalis</i> (Günther, 1858)	LC	20 (9.20%)	18 (9.80%)	2 (5.90%)
<b>Phrynobatrachidae</b>						
	<i>Phrynobatrachus</i>					
		<i>P. plicatus</i> (Günther, 1858)	LC	57 (26.20%)	57 (31.00%)	0 (0.00%)
		<i>P. auritus</i> (Boulenger, 1900)	LC	9 (4.10%)	9 (4.90%)	0 (0.00%)
<b>Ptychadenidae</b>						
	<i>Ptychadena</i>					
		<i>P. bibronii</i> (Hallowell, 1845)	LC	3 (1.40%)	3 (1.60%)	0 (0.00%)
		<i>P. longirostris</i> (Peters, 1870)	LC	45 (20.60%)	45 (24.50%)	0 (0.00%)
		<i>P. oxyrhynchus</i> (Smith, 1849)	LC	19 (8.70%)	19 (10.30%)	0 (0.00%)
<b>TOTAL</b>				<b>218</b>	<b>184</b>	<b>34</b>

Note: IUCN Status: LC= Least Concern, CE= Critically Endangered; Values in the bracket represent the relative abundance.