

Modeling the past and contemporary habitat suitability and distribution of the Levantine viper *Macrovipera lebetinus* (Linnaeus, 1758) (Ophidia: Viperidae)

Rasoul Karamiani*, Maryamalsadat Hosseini

Department of Biology, Faculty of Science, Razi University, Kermanshah, Iran

* Email: r.karamiani@razi.ac.ir

Received: 01 August 2023 / Revised: 08 September 2023 / Accepted: 09 September 2023/ Published online: 10 September 2023.

How to cite: Karamiani, R., Hosseini, M. (2024). Modeling the past and contemporary habitat suitability and distribution of the Levantine viper *Macrovipera lebetinus* (Linnaeus, 1758) (Ophidia: Viperidae), Journal of Wildlife and Biodiversity, 8(1), 39-53. DOI: <https://doi.org/10.5281/zenodo.10162258>

Abstract

Evaluating the climate changes in the past and contemporary times, and the impact of those changes on the distribution range of the species has attracted research interest. The venomous snakes of the genus *Macrovipera* consisting of the recognized species *M. lebetinus* and *M. razii* are documented from Iran. In the study, we modeled the potential distribution areas and determined the suitable habitats in the past (the Last Interglacial [LIG], and mid-Holocene [MH]), and their present distribution for the Levantine viper *M. lebetinus* by MaxEnt. Models of the species indicated good fit operation by the average high area under the curve (AUC) values (LIG = 0.979 ± 0.008 , MH = 0.968 ± 0.028 , Contemporary = 0.933 ± 0.036). Three important climate variables had significant contributions to the simulation of the LIG model distribution of *M. lebetinus* as mean temperature of the driest quarter of the year (43.8%), isothermality (26.9%), and mean temperature of the wettest quarter of the year (17.9%); the maximum temperature of the warmest month (36.4%), mean temperature of the wettest quarter of the year (25.2%), and isothermality (21.2%) variables had significant contributions to the contemporary time and the MH model distribution the Levantine viper, respectively; Two important climate variables had immense contributions to the prediction of the contemporary model distribution *M. lebetinus* as seasonal and annual precipitation (32.9%), and a topographic variable as slope (31.5%). Because it seems that they have sensitivity to temperature and precipitation levels of the seasons. The MH and the LGM models indicated a larger suitable area than the contemporary distribution, and it is concluded that these variables form a natural barrier for species dispersion.

Keywords: Blunt-nosed viper, Climate condition, Last interglacial, mid-Holocene

Introduction

The portentous response of species to the continuous climate changes may be incessantly *in situ* at their tolerance limits or changing ranges in the regions where the climate is within the species

tolerance limits, and ultimately extinction (Davis et al., 2005). The glaciations were interrupted by interglacial periods (Ray, 1992), the climate warming caused the forests to be revived in the areas where once supported tundra vegetation (Sillero & Carretero, 2012), and climate of the Last Interglacial period (LIG: 150,000–115,000 years BP) was warmer than the present time with an increased temperature gradient (~3-5 °C) in polar regions towards lower latitudes and caused the sea-level to rise (approximately 4 to 6 m) and the ice sheets to reduce (Pickarski, 2014; Nikolova et al., 2013). During the mid-Holocene (6k), summer temperatures were warmer than the present time in the high latitudes of the Northern Hemisphere, but northern Africa, Arabia, and southern Asia (Iran and surrounding regions) underwent the conditions that the precipitation and humidity increased compared to the present time and these conditions resulted in both African and Asian monsoons (Wanner et al., 2008; Jones et al., 2013). Species distribution models (SDMs) can be capitalised on to investigate the impingement of climate changes on the distributions and abundances of species (Thomas et al., 2004); analysing SDMs can assist in conserving management planning (Graham et al., 2004, Farahani and Asgharzadeh, 2023), understanding theoretical research on ecological processes (Phillips et al., 2009), determining biogeographical patterns, predicting potential distribution (Ananjeva et al., 2006), appraising possible future changes in the diversity (Ramirez-Villegas et al., 2014) and predicting potential distribution and habitat suitability for species (Karamiani et al., 2018). MaxEnt is a general approach for characterising the probable distributions from small sample sizes (Elith et al., 2006; Phillips et al., 2006). MaxEnt estimates the probable distribution of maximum entropy (i.e. closest to uniform) based on environmental variables dispersed over the areas to be surveyed (Pearson et al., 2007; Kaliontzopoulou et al., 2008).

Viperidae family (with subfamilies Azemiopinae: Fea's vipers, Crotalinae: Pit vipers, and Viperinae: True vipers) has more than 9.4% of all the living genera and species of the snakes (Uetz et al., 2023). The venomous snakes of the genus *Macrovipera* Reuss, 1927 (Type species: *Coluber lebetinus* Linnaeus, 1758) were distinguished and described from the genus *Vipera* Laurenti, 1768 by Herrmann et al., (1992) in order to include the large North African and Eurasian vipers (Cattaneo, 2020). The genus *Macrovipera* consists of two species as follows: *M. lebetinus* (Linnaeus, 1758), and *M. razii* Oraie, Rastegar-Pouyani, Khosrovani, Moradi, Akbari, Sehhatisabet, Shafiei, Stümpel & Joger, 2018 which were geographically distributed in Morocco, Algeria, Tunisia, Cyprus, Syria, Lebanon, Jordan, Turkey, Iraq, Yemen (recorded by Scortecci 1932), Iran, Afghanistan, Pakistan, India (Jammu and Kashmir) (Leviton et al., 1992; Khan 2004; Stümpel & Joger, 2009; Uetz et al., 2023). The Levantine viper *M. lebetinus* is known with six recognized and valid subspecies: The

nominal subspecies Cypriot blunt-nosed viper *M. l. lebetinus* (Linnaeus 1758) seems to be endemic to Cyprus; the Chernov's viper *M. l. cernovi* (Chikin and Szczerbak 1992) in Iran (Kerman, Khorasan Razavi, North Khorasan and Semnan Provinces), Afghanistan, Pakistan, Turkmenistan, Kyrgyzstan, Uzbekistan, India (Kashmir); the Levant blunt-nosed viper *M. l. obtusa* (Dwigubsky 1832) in Lebanon, Jordan, Iraq, Syria, northeast Turkey, Transcaucasia, Iran (West Azarbaijan, East Azarbaijan, Ardabil, Qazvin, Gilan, Mazandaran, Golestan, Alborz, Tehran, Semnan, Qom, Markazi, Zanjan, Hamedan, Kurdistan, Kermanshah, Lorestan, Ilam, Khuzestan, Kohgiluyeh Va Boyer Ahmad, Fars and Esfahan Provinces), Afghanistan, Pakistan; the Milos viper *M. l. schweizeri* (formally *Viper schweizeri* Werner 1935) in Greece; *M. l. transmediterranea* (Nilson and Andr n 1988) in Algeria, Tunisia; and the Turan blunt-nosed viper *M. l. turanica* (Chernov 1939) in Tajikistan, Uzbekistan, Kyrgyzstan (Ananjeva et al., 2006; Jestrzemiński & Kuzyakova, 2018; Uetez et al., 2022). According to the successful results of the hybridisation of the subspecies *schweizeri*, *lebetina*, *obtusa* and *turanica*, they have not raised the full species, because interbreeding has occurred between them (Kamelin et al., 1997).

In addition to the above-mentioned subspecies, there is also a challenging subspecies *M. l. peilei* (Murray, 1892), in Pakistan, which was rejected by the group of researchers (Golay et al., 1993; McDiarmid et al., 1999) and it is categorised as a synonym of *obtuse*, but according to Sindaco et al., (2013) *peilei* is synonymous with *M. lebetina cernovi*. Cattaneo (2020) investigated five species of *Macrovipera*: *lebetinus*, *razii*, *schweizeri*, *mauritanica* and *deserti* that *schweizeri*, *mauritanica* and *deserti* are considered subspecies of *lebetinus*, and Lenk et al., (2001) subcategorised *mauritanica* and *deserti* into the distinct genus *Daboia*, while other researchers are currently invalid (*euphratica*, *siphnensis*). The genus *Macrovipera* in the molecular phylogenetic aspect is a sister taxon of the mountain viper *Montivipera* Nilson, Tuniyev, Andr n, Orlov, Joger and Herrmann, 1999 (Lenk et al., 2001, Garrigues et al., 2005; W ster et al., 2008; St mpel & Joger, 2009; Pyron et al., 2013; Oraie et al., 2018). The general aim of this work is: 1) to identify model potential areas of distribution during three periods of the past: Last Interglacial (LIG: ~120,000–140,000 years BP) mid-Holocene (MH: ~6,000 years BP), 2) to describe present distribution, suitable habitat, and understand the biogeographical patterns of *M. lebetinus* in World.

Martial and methods

Study area and presence records

The Levantine viper *M. lebetinus* (Fig. 1) can usually be found in the mountainous areas of Alborz

and Zagros and their adjacent areas, even in rural residential areas. This species is nocturnal and not very aggressive snake and it can also be seen in the early morning or at dusk. We collected the data concerning the species occurrence based on systematic biological survey by walking randomly through the habitat from 06:00 to 08:00 AM and 18.00-22:00 PM to evening (much of the activity time of species) during the spring to summer 2013 and 2021. We used localities mentioned in previous studies (e. g. Latifi, 1991; Disi et al., 2001; Khan 2002; Hraoui-Bloquet et al., 2002; Jablonski & Masroor, 2020). We gathered the data related to the distribution of *M. lebetinus* specimens collected in rodents' nests, groove of rocks or around bushes of oak forest in the Zagros and Kopet Dag Mountains, near a poultry farm (Javanrod, Kurdistan Province) (Fig. 2) and recorded the exact location by using the Global Positioning System (GPS), and localities from the Global Biodiversity Information Facility (GBIF: <https://www.gbif.org>) website (GBIF 2022) were also included. In total, 140 records were gathered and used in this study.



Fig. 1. Alive adult specimen of *Macrovipera lebetinus* in Zoology laboratory, Razi University (A); the camouflage of Levantine viper in its habitat (Tandoureh Protected Area northeast, near the city of Dargaz and the Turkmenistan border) (B and C).




Fig. 2. Landscape of habitat of *Macrovipera lebetinus* in Pouyan Region, Kermanshah Province (A); Malle Punjab, Ilam Province (B); Bloran Region, Lorestan Province (C), and Tandoureh Protected Area, Khorasan Razavi Province (D).

Data set and analysis

We carried out the Maximum Entropy modeling (MaxEnt) using the software MaxEnt, 3.3.3e (Phillips et al., 2006) for surveying species geographic distributions and the data to test samples. In this study, 19 bioclimatic variables and two topographical variables were used for the contemporary time, and also 19 bioclimatic variables were used in the past time (LIG, and MH) in the species-related parts of the world (Eurasian and North African) (Hijmans et al., 2005; Otto-Bliesner et al., 2006; www.worldclim.org) (see the Appendix). To recognise the correlation ratios between variables and present records, Openmodeller (V. 1.0.7) (de Souza Muñoz et al., 2011), was used. Then, we used SPSS IBM (version 22) for Pearson's correlation coefficient (Elith et al., 2006). We selected variables with a Pearson correlation lower than 0.75 to select the variables that are ecologically important for species separation according to our observations, in order for the habitat to be described.

We applied MaxEnt software and the analysis was repeated 15 times, yielding the best model for the studied species. Twenty-five percent of the data was used as test data, and the remaining percent of data (75 percent) was considered as training data. MaxEnt provides the distribution models by the receiver operating characteristic (ROC) plots, ROC curves plot true-positive rate against false-positive rates (Kaliontzopoulou et al., 2008). A value of 0.5–0.7 related to the area under the curve (AUC) indicates that the result is a stochastic prediction (Gallien et al., 2012), and values of 0.7–0.9 suggest an efficient model, the values more than 0.9 indicate high accuracy (Manel et al., 2001). We used DIVA-GIS 7.3.0.1 software for predicting the maps and a logistic output of present records with a suitability range showing the values of zero (unsuitable habitat) to one (the best suitable habitat) (Hijmans et al., 2001).

Results

The habitat suitability and distribution of the Levantine viper *M. lebetinus* were best described by a combination of bioclimatic and geographic variables suggested by a high value of the area under the curve (AUC) (LIG = 0.979 ± 0.008 , MH = 0.968 ± 0.028 , and contemporary period = 0.933 ± 0.036). The models showed a very significant match and closely fitted the presence of the species recorded in the studied areas. The contribution of different variables to different time periods is shown in Table 1. In the LIG model, the Mean temperature of the driest (19.7 ± 8.6 °C), and wettest (9.8 ± 5.3 °C) quarter of the year revealed more than half of the variance (61.7%); Therefore, temperature modality was the most important factor for the simulated distribution model during the LIG periods (Fig. 3A). In the MH model, maximum temperature of the warmest month (45.4 °C), and mean temperature of the wettest quarter of the year (10.2 °C) showed 61.6% of the variance. Also, the temperature was the most important factor for the simulated distribution models during the MH periods (Fig. 3B). In the contemporary model, seasonal and annual precipitation justified 32.9% and the slope accounts for 31.5% of the variance, where mean precipitation was 67-357 mm with slope 89° (Fig. 3C). The model for *M. lebetinus* predicated the occurrence range of the species in rocky hills, and mountain sides with an elevation of 3 to 2640 m above sea level (a.s.l.).

Table 1. Relative importance of variables (in percentages) in the last interglacial (LIG, ~120 ka), mid-Holocene (MH, ~6 ka), and the present period used in MaxEnt model for the Levantine viper *M. lebetinus*.

Variable	Description of variables	LIG	MH	Present time
Bio9	Mean temperature of the driest quarter of the year	43.8	-	-
Bio5	Maximum temperature of the warmest month	-	36.4	-
Bio3	Isothermality [(BIO2 / BIO7) * 100]	26.9	21.2	11.5
Bio8	Mean temperature of the wettest quarter of the year	17.9	25.2	-
Bio15	Precipitation seasonality (standard deviation / mean)	9.4	12.9	16.5
Bio2	Mean diurnal range [mean of monthly (max temp – min temp)]	2.1	4.3	-
Slope	Slope	-	-	31.5
Bio12	Precipitation seasonality Annual precipitation	-	-	16.4
Bio4	Temperature seasonality (standard deviation * 100)	-	-	15.9
Altitude	Altitude	-	-	8.2




Fig. 3 Distribution map of *Macrovipera lebetina* and much of their potential distribution pattern in the region during: A) the last interglacial (120 ka), B) the mid-Holocene (6 ka), C) currently. Af. Afghanistan, Al. Albania, Alg. Algeria, Az. Azarbaijan, Chi. China, Cy. Cyprus, Eg. Egypt, Fr. France, Gr. Greece, In. India, Ira. Iraq, Ir. Iran, It. Italy, Li. Libya, Ka. Kazakhstan, Ky. Kyrgyzstan, Mo. Morocco, Pa. Pakistan, Po. Portugal, Sp. Spain, Sy. Syria, Ta. Tajikistan, Tur. Turkey, Tu. Tunisia, Turk. Turkmenistan, Uz. Uzbekistan.

Discussion

Results of the present study verify the known distribution of the Levantine viper based on climatic conditions. The regions in the vicinity of the Mediterranean Sea, western Asia (the Middle East and the Caucasus), middle Asia (the plains and hills lying between the Caspian Sea to the west of China), south Asia (Afghanistan, Pakistan, and India) had a wide variety of area suitability for the presence of the Levantine viper during LIG, MH, contemporary periods. Climate change and habitat destruction (fragmentation and loss) are considered threats to global biodiversity (Travis, 2003; Opdam & Wascher, 2004). Climate changes followed by habitat destruction are the primary causes of the decline of reptiles globally and trigger the local extinction in many parts of the world. The habitat suitability such as northern Uzbekistan and Turkmenistan, southern and central Iraq, north Egypt and northeastern Libya have been lost for the Levantine viper over time; when habitat is lost, the populations of species are also likely to decline and disappear (e.g. Bender et al., 1998; Andr n, 1999; Mantyka-pringle et al., 2012). In the LIG, MH, contemporary periods, habitats in northern India and western Jordan had remained as patches with relatively low suitability (18 to 36%); the process of habitat fragmentation with increased number, isolation of patches and small populations increase the risk of extinction (Young et al., 1996; Mantyka-pringle et al., 2012). However, according to the models, the habitats of the west of the Caspian Sea have become more suitable for the Levant viper, and habitats such as southern Italy, Spain and Portugal were suitable in the three time periods, but until now there have been no report of the Levantine viper in the above-mentioned regions, which is possibly due to lack of access to those habitats. The Levantine viper inhabit in the very diverse desert and montane-steppe biotopes (Ananjeva et al., 2006) has made the viper to be able to tolerate climate change imposed on its habitats and continue to survive. Most records of the species occur in rock and dry foothills, on mountain slopes with sparse annual grasses, and thorny bushes, rocky ravines with small streams and springs, in pistachio and oak woodlands, on precipices in river valleys, on the banks of irrigation channels (Leviton et al., 1992; Ananjeva et al., 2006). The results of present study are consistent with all the previous records (e.g. Disi, et al., 2001; Ananjeva et al., 2006; Cattaneo, 2020) except the record of Yemen (a specimen collected in 1932 by Scortecci with uncertain locality), which was not suitable for inhabiting the Levantine viper during the LIG, MH, and contemporary periods. The results showed that the distribution of the Levantine viper is limited by different climatic parameters. Model distribution ranges during the MH to the present time indicated a gradual decline of habitat suitability in Morocco to northern India. During spring and summer, the increased

temperature and rainfall fluctuations over the regions show a clear decrease from the MH to contemporary time which can be elucidated by the changes in solar insolation due to the varying orbital force (Fallah et al., 2017). Park et al., (2019) reconstructed the annual mean Northern Hemisphere warming and Arctic amplification in response to Mid-Holocene insolation and also simulated the pronounced summer warming anomalies that persisted into winter. Terral & Mengüal (1999) suggested annual temperatures between 1.5°C and 3.5°C for southern Europe and the areas around the Mediterranean Sea (Molnar & Rajagopalan, 2020), which were cooler and wetter than the present time. Therefore, the MH had suitable climate conditions for the distribution of the Levantine viper from areas around the Mediterranean Sea to north India and west Kyrgyzstan. We think that temperature and precipitation, especially during spring and summer, are the most effective factors determining the habitat suitability for *M. lebetina*.

In addition to habitat change, fragmentation and decreasing connectivity between suitable habitats (small and isolated patches) lead to the interruption of gene flow among populations and, ultimately an increased extinction risk (Prugh et al., 2008; Fischer & Lindenmayer 2007; Ferraz et al., 2007; Spiesman et al., 2018). Three patterns are simulated for climate changes in our modelling: first, some suitable habitats almost disappear in the areas of climate change, as in northern Uzbekistan and Turkmenistan, southern and central Iraq, north Egypt and northeastern Libya; second, the vastness of suitable habitats reduces, as in Iran, Turkey, Syria, and Jordan; third, the suitability of some habitats increases, as in Azerbaijan, Armenia, and Dagestan. Consequently, climate change can cause the disappearance or reduction as well as the expansion of suitable habitats (Parmesan 2006; Bonino et al., 2015; Kim et al., 2020; Karamiani & Rastegar-Pouyani, 2021). Therefore, these regions might function as potential refuges for conservation of the species under the climate changes in the future.

Conclusion

The present study was carried out with the aim of modeling the potential distribution areas and determined the suitable habitats in the past (the Last Interglacial, and mid-Holocene), and current distribution of the Levantine viper by MaxEnt. Based on the results of this study can be firstly understand the decreasing trend of habitats with the effect of climate change from the past to the present, and secondly, to identify the range of distribution and habitats of the Levantine viper for can help to reduce the possibility of snakebite.

Acknowledgements

We wish to thank Eskandar Rastegar-Pouyani, Seyyed Saeed Hosseinian Yousefkhani, Ali Permanoon, Erfan Permanoon, Akbar Fattahi, Hossain Nabizadeh and Morad Besharati for assisting us with field work.

References

- Ananjeva, N.B., Orlov, N.L., Khalikov, R.G., Darevsky, I.S., S.A., Ryabov, I.S., & Barabanov, A.V. (2006). The reptiles of northern Eurasia: taxonomic diversity, distribution, conservation status (No. 47). Pensoft Publishers.
- Andr n, H. (1999). Habitat fragmentation, the random sample hypothesis and critical thresholds. *Oikos*, 84 (2), 306–308.
- Bender, D.J., Contreras, T.A., & Fahrig, L. (1998). Habitat loss and population decline: a meta-analysis of the patch size effect. *Ecology*, 79(2), 517–533.
- Bonino, M.F., Moreno Azocar, D.L., Schulte, J.A., & Cruz, F.B. (2015). Climate change and lizards: changing species' geographic ranges in Patagonia. *Regional Environmental Change*, 15 (6), 1121–1132.
- Cattaneo, A. (2020). *Macrovipera schweizeri* (Werner, 1935): comparison of the populations of Milos and Sifnos islands (SW Cyclades) (Serpentes Viperidae). *Naturalista sicil*, 44, 127–54.
- Davis, M.B., Shaw, R.G., & Etterson, J.R. (2005). Evolutionary responses to changing climate. *Ecology*, 86(7), 1704 – 1714.
- De Souza Mu oz, M.E., De Giovanni, R., de Siqueira, M.F., Sutton, T., Brewer, P., Pereira, R.S., Canhos, D.A.L., & Canhos, V.P. (2011). openModeller: a generic approach to species' potential distribution modelling. *GeoInformatica* 15 (1), 111–135.
- Disi, A.M., Modry, D., Necas, P., & Rifai, L. (2001). Amphibians and Reptiles of the Hashemite Kingdom of Jordan: An Atlas and Field Guide Edition Chimaira. Frankfurt am Main, 408 pp.
- Elith, J., Graham, C.H., Anderson, R. P., Dud k, M., Ferrier, S., Guisan, A., Hijmans, R.J., Huettmann, F., Leathwick, J.R., Lehmann, A., Li, J., Lohmann, L.G., Loiselle, B.A., Manion, G., Moritz, C., Nakamura, M., Nakazawa, Y., Overton, J.M., Peterson, A.T., Phillips, S.J., Richardson, K.S., Scachetti- Pereira, R., Schapire, R.E., Soberon, J., Williams, S., Wisz, M.S., & Zimmermann, N.E. (2006). Novel methods improve prediction of species' distributions from occurrence data. *Ecography*, 29(2), 129–151.
- Fallah, B., Sodoudi, S., Russo, E., Kirchner, I., & Cubasch, U. (2017). Towards modeling the regional rainfall changes over Iran due to the climate forcing of the past 6000 years. *Quaternary International*, 429, 119–128.
- Farahani, R., & Asgharzadeh, A. (2023). Golestan National Park's ecosystem services, a case with brown bears habitat suitability. *Scientific Reports in Life Sciences*, 4(1), 1–7. <https://doi.org/10.5281/zenodo.7632690>
- Ferraz, G., Nichols, J.D., Hines, J.E., Stouffer, P.C., Bierregaard Jr, R.O., & Lovejoy, T.E. (2007). A large-scale deforestation experiment: effects of patch area and isolation on Amazon birds. *science*, 315(5809), 238–241.
- Fischer, J., & Lindenmayer, D.B. (2007). Landscape modification and habitat fragmentation: a synthesis. *Global ecology and biogeography*, 16(3), 226565-280.
- Gallien, L., Douzet, R., Pratte, S., Zimmermann, N.E., & Thuiller, W. (2012). Invasive species distribution models—how violating the equilibrium assumption can create new insights. *Global Ecology and Biogeography*, 21(11), 1126–1136.

- Garrigues, T., Dauga, C., Ferquel, E., Choumet, V., & Failloux, A.B. (2005). Molecular phylogeny of *Vipera Laurenti*, 1768 and the related genera *Macrovipera* (Reuss, 1927) and *Daboia* (Gray, 1842), with comments about neurotoxic *Vipera aspis aspis* populations. *Molecular Phylogenetics and Evolution*, 35(1), 35–47.
- Golay, P., Smith, H.M., Broadley, J., Dixon, J.R., McCarthy, C., Rage, J.C., & Schätti, B., & Toriba, M. (1993). Endoglyphs and other major venomous snakes of the world: a checklist. Azemiops S. A., Geneva, Switzerland.
- Graham, C.H., Ferrier, S., Huettman, F., Moritz, C., & Peterson, A.T. (2004). New developments in museum-based informatics and applications in biodiversity analysis. *Trends in ecology & evolution*, 19(9), 497–503.
- Herrmann, H.W., Joger, U., & Nilson, G. (1992). Phylogeny and systematics of viperine snakes. III: resurrection of the genus *Macrovipera* (Reuss, 1927) as suggested by biochemical evidence. *Amphibia-Reptilia*, 13(4), 375–392.
- Hijmans, R.J., Guarino, L., Cruz, M., & Rojas, E. (2001). Computer tools for spatial analysis of plant genetic resources data: 1. DIVA-GIS. *Plant genetic resources newsletter*, 15–19.
- Hraoui-Bloquet, S., Sadek, R.A., Sindaco, R., & Venchi, A. (2002). The herpetofauna of Lebanon: new data on distribution. *Zoology in the Middle East*, 27(1), 35–46.
- <http://www.gbif.org>. Occurrence Download <https://doi.org/10.15468/dl.ywygqv> [Accessed: 08 March 2022].
- Jablonski, D., & Masroor, R. (2020). *Macrovipera lebetinus* in Pakistan. *Herpetological Bulletin*, 153, 44–45.
- Jestrzanski, D., & Kuzyakova, I. (20198). Morphometric characteristics and seasonal proximity to water of the Cypriot bluntnosed viper *Macrovipera lebetina lebetina* (Linnaeus, 1758). *Journal of Venomous Animals and Toxins including Tropical Diseases*, 24.
- Jones, M., Djamali, M., Stevens, L., Heyvaert, V., Askari, H., Noorollahi, D.W., & Weeks, L. (2013). Mid-Holocene environmental and climatic change in Iran. *Ancient Iran and its neighbours: local developments and long-range interactions in the 4th millennium BC*, British Institute for Persian Studies and Oxbow Books, Oxford, UK, 25–34.
- Kaliontzopoulou, A., Brito, J.C., Carretero, M.A., Larbes, S., & Harris, D.J. (2008). Modelling the partially unknown distribution of wall lizards (*Podarcis*) in North Africa: ecological affinities, potential areas of occurrence, and methodological constraints. *Canadian Journal of Zoology*, 86(9), 992–1001.
- Kamelin, E.R., Lukin, Y.A., & Milto, K.D. (1997). Hybridization of *Vipera schweizeri* (Werner, 1935) and *Vipera lebetina obtusa*, Dvigubsky 1832. *Russian Journal of Herpetology*, 4(1), 75–78.
- Karamiani, R., & Rastegar-Pouyani, N. (2021). The effect of climate change on habitat suitability and a distribution model of the Iranian fat-tailed gecko, *Eublepharis angramainyu* Anderson and Leviton, 1966 (Sauria: Eublepharidae) since the last interglacial to 2050. *Zoology and Ecology*, 31, 24–32.
- Karamiani, R., Rastegar-Pouyani, N., & Rastegar-Pouyani, E. (2018). Modeling the past and current distribution and habitat suitability for *Ablepharus grayanus* and *A. pannonicus* (Sauria: Scincidae). *Asian Herpetological Research*, 9(1), 56–64.
- Khan, M.S. (2002). *A Guide to the Snakes of Pakistan*. Frankfurt am Main: Edition Chimaira.
- Kim, D.I., Park, I.K., Bae, S.Y., Fong, J.J., Zhang, Y.P., Li, S.R., Ota, J. Kim, S. & Park, D. (2020). Prediction of present and future distribution of the Schlegel's Japanese gecko (*Gekko japonicus*) using MaxEnt modeling. *Journal of Ecology and Environment*, 44(1), 1–8.
- Latifi, M. (1991). *The Snakes of Iran*. English edition. Contributions to Herpetology, vol. 7. Ohio, USA: SSAR Oxford, 159 pp.
- Lenk, P., Kalyabina, S., Wink, M., & Joger, U. (2001). Evolutionary relationships among the true vipers (Reptilia: Viperidae) inferred from mitochondrial DNA sequences. *Molecular phylogenetics and*

- evolution, 19(1), 94–104.
- Leviton, A.E., Anderson, S.C., Adler, K.A., & Minton, S.A. (1992). Handbook to Middle East Amphibians and Reptiles. Oxford, Ohio Vii + 252 pp.
- Mantyka-pringle, C.S., Martin, T.G., & Rhodes, J.R. (2012). Interactions between climate and habitat loss effects on biodiversity: a systematic review and meta-analysis. *Global Change Biology*, 18(4), 1239–1252.
- McDiarmid, R. W., Campbell, J. A., & Touré, T. (1999). Snake species of the world: A taxonomic and geographic reference. Herpetologists' League, Washington, DC.
- Molnar, P., & Rajagopalan, B. (2020). Mid-Holocene Sahara-Sahel precipitation from the vantage of present-day climate. *Geophysical Research Letters*, 47(16), e2020GL088171.
- Nikolova, I., Yin, Q., Berger, A., Singh, U.K., & Karami, M.P. (2013). The last interglacial (Eemian) climate simulated by LOVECLIM and CCSM3. *Climate of the Past*, 9(4), 1789–1806.
- Opdam, P., & Wascher, D. (2004). Climate change meets habitat fragmentation: linking landscape and biogeographical scale levels in research and conservation. *Biological conservation*, 117(3), 285–297.
- Oraie, H., Rastegar-Pouyani, E., Khosravani, A., Moradi, N., Akbari, A., Sehhatiasabet, M.E., Shafiei, S., Stümpel, N. & Joger, U. (2018). Molecular and morphological analyses have revealed a new species of blunt-nosed viper of the genus *Macrovipera* in Iran. *Salamandra*, 54(4), 233–248.
- Park, H.S., Kim, S.J., Stewart, A.L., Son, S.W., & Seo, K.H. (2019). Mid-Holocene Northern Hemisphere warming driven by Arctic amplification. *Science advances*, 5(12), eaax8203.
- Parmesan, C. (2006). Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology Evolution and Systematics* 37, 637–669.
- Pearson, R.G., Raxworthy, C.J., Nakamura, M., & Townsend Peterson, A. (2007). Predicting species distributions from small numbers of occurrence records: a test case using cryptic geckos in Madagascar. *Journal of biogeography*, 34(1), 102–117.
- Phillips, S.J., Anderson, R.P., & Schapire, R.E. (2006). Maximum entropy modeling of species geographic distributions. *Ecological modelling*, 190(3–4), 231–259.
- Phillips, S.J., Dudík, M., Elith, J., Graham, C.H., Lehmann, A., Leathwick, J., & Ferrier, S. (2009). Sample selection bias and presence-only distribution models: implications for background and pseudo-absence data. *Ecological applications*, 19(1), 181–197.
- Pickarski, N. (2014). Vegetation and climate history during the last glacial–interglacial cycle at Lake Van, eastern Anatolia. Thesis. Universitäts- und Landesbibliothek Bonn.
- Prugh, L.R., Hodges, K.E., Sinclair, A.R., & Brashares, J.S. (2008). Effect of habitat area and isolation on fragmented animal populations. *Proceedings of the National Academy of Sciences*, 105(52), 20770–20775.
- Pyron, R.A., Burbrink, F.T., & Wiens, J.J. (2013). A phylogeny and revised classification of Squamata, including 4161 species of lizards and snakes. *BMC evolutionary biology*, 13, 1–54.
- Ramirez-Villegas, J., Cuesta, F., Devenish, C., Peralvo, M., Jarvis, A., & Arnillas, C.A. (2014). Using species distributions models for designing conservation strategies of Tropical Andean biodiversity under climate change. *Journal for Nature Conservation*, 22(5), 391–404.
- Ray, L. L. (1992). The Great Ice Age. US Department of the Interior, US Geological Survey.
- Sillero, N., & Carretero, M.A. (2013). Modelling the past and future distribution of contracting species. The Iberian lizard *Podarcis carbonelli* (Squamata: Lacertidae) as a case study. *Zoologischer Anzeiger-A Journal of Comparative Zoology*, 252(3), 289–298.
- Sindaco, R., Venchi, A. & Grieco, C. (2013). The Reptiles of the Western Palearctic, Volume 2: Annotated Checklist and Distributional Atlas of the Snakes of Europe, North Africa, Middle East and Central Asia, with an Update to. *Monografie della Societas Herpetologica Italica*.

- Spiesman, B.J., Stapper, A.P., & Inouye, B.D. (2018). Patch size, isolation, and matrix effects on biodiversity and ecosystem functioning in a landscape microcosm. *Ecosphere*, 9(3), e02173.
- Stümpel, N., & Joger, U. (2009). Recent advances in phylogeny and taxonomy of Near and Middle Eastern Vipers—an update. *ZooKeys*, 31, 179–191.
- Terral, J.F., & Mengüal, X. (1999). Reconstruction of Holocene climate in southern France and eastern Spain using quantitative anatomy of olive wood and archaeological charcoal. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 153(1–4), 71–92.
- Thomas, C.D., Cameron, A., Green, R.E., Bakkenes, M., Beaumont, L.J., Collingham, Y.C., ... & Williams, S. E. (2004). Extinction risk from climate change. *Nature*, 427(6970), 145–148.
- Travis, J.M.J. (2003). Climate change and habitat destruction: a deadly anthropogenic cocktail. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 270(1514), 467–473.
- Uetz, P., Freed, P., Aguilar, R., Reyes, F., & Hošek, J. (2023). The Reptile Database, URL: <http://www.reptile-database.org> (accessed on June 12, 2023).
- Wanner, H., Beer, J., Bütikofer, J., Crowley, T.J., Cubasch, U., Flückiger, J., ... & Widmann, M. (2008). Mid-to Late Holocene climate change: an overview. *Quaternary Science Reviews*, 27(19–20), 1791–1828.
- Wüster, W., Peppin, L., Pook, C.E., & Walker, D.E. (2008). A nesting of vipers: phylogeny and historical biogeography of the Viperidae (Squamata: Serpentes). *Molecular Phylogenetics and Evolution*, 49(2), 445–459.
- Young, A., Boyle, T., & Brown, T. (1996). The population genetic consequences of habitat fragmentation for plants. *Trends in ecology & evolution*, 11(10), 413–418.

Appendix

Table 1. Climatic variables are used to elaborate the models (www.worldclim.org).

Characters	Definition
Altitude	Altitude
BIO1	Annual mean temperature
BIO2	Mean diurnal range [mean of monthly (max temp - min temp)]
BIO3	Isothermality [(BIO2 / BIO7) * 100]
BIO4	Temperature seasonality (standard deviation * 100)
BIO5	Maximum temperature of the warmest month
BIO6	Minimum temperature of the coldest month
BIO7	Temperature annual range (BIO5 - BIO6)
BIO8	Mean temperature of the wettest quarter of the year
BIO9	Mean temperature of the driest quarter of the year
BIO10	Mean temperature of the warmest quarter of the year
BIO11	Mean temperature of the coldest quarter of the year
BIO12	Annual precipitation
BIO13	Precipitation of the wettest month
BIO14	Precipitation of the driest month
BIO15	Precipitation seasonality (standard deviation / mean)
BIO16	Precipitation of the wettest quarter of the year
BIO17	Precipitation of the driest quarter of the year
BIO18	Precipitation of the warmest quarter of the year
BIO19	Precipitation of the coldest quarter of the year
Slope	Slope